of the year (Andrewartha & Barker, 1969). In comparison, the Bennett's wallaby in Tasmania (present study) appears to show no less synchrony but with births occurring about two weeks later. In both species, births during the period of seasonal quiescence are uncommon (Andrewartha & Barker, 1969; Catt, 1977; Merchant & Calaby, 1981) but can occur in a very low proportion of animals (tammar: Hinds & den Ottolander, 1983; Bennett's wallaby: present study). Survival of such young may be poor as peak lactation and pouch emergence would occur during the winter.

I thank Professor Mike Stoddart and colleagues at the Department of Zoology, University of Tasmania, Hobart for the provision of moral support and departmental facilities during my visit to Tasmania. The field trip to Tasmania was funded by a Wellcome-Ramaciotti Research Travel Grant and a grant from the Nuffield Foundation.

REFERENCES

- Andrewartha, H. G. & Barker, S. (1969). Introduction to a study of the ecology of the Kangaroo Island wallaby Protemnodon eugenii (Desmarest) within Flinders Chase, Kangaroo Island, South Australia. Trans. R. Soc. S. Aust. 93: 127-133.
- Catt, D. C. (1977). The breeding biology of Bennett's wallaby (Macropus rufogriseus rufogriseus) in South Canterbury, New Zealand. N.Z. J. Zool. 4: 401-411.
- Fleming, D., Cinderey, R. N. & Hearn, J. P. (1983). The reproductive biology of Bennett's wallaby (*Macropus rufogriseus* rufogriseus) ranging free at Whipsnade Park. J. Zool., Lond. 201: 283-291.
- Hinds, L. A. & den Ottolander, R. C. (1983). Effect of changing photoperiod on peripheral plasma prolactin and progesterone concentrations in the tammar wallaby (*Macropus eugenii*). J. Reprod. Fert. 69: 631-639.
- Merchant, J. C. & Calaby, J. H. (1981). Reproductive biology of the Red-necked wallaby (Macropus rufogriseus banksianus) and Bennett's wallaby (M. r. rufogriseus) in captivity. J. Zool., Lond. 194: 203-217.
- Poole, W. E., Carpenter, S. M. & Wood, T. J. (1982). Growth of Grey kangaroos and the reliability of age determination from body measurements. I. The Eastern grey kangaroo, *Macropus giganteus*. Aust. Wildl. Res. 9: 9-20.
- Tyndale-Biscoe, C. H., Hearn, J. P. & Renfree, M. B. (1974). Control of reproduction in macropodid marsupials. J. Endocr. 63: 589-614.

J. Zool., Lond. (1989) 218, 339-343

Reproductive strategies in female Spanish ibex (Capra pyrenaica)

Paulino Fandos, Estación Biológica de Doñana, Unidad de Ecologia, Pabellón del Perú. Avda. Maria Luisa s/n, 41013 Sevilla, España*

The reproductive performance of two populations of the Spanish ibex (Capra pyrenaica) has been analysed by direct observations. One population was living in natural conditions, the other in semi-captivity where a daily food supplement was given. The most important differences were in quantitative aspects of reproduction, such as the age of sexual maturity and the litter size. This difference in the reproductive strategy can be explained either by the availability of food or the severity of winter.

^{*} Present address: Museo Nacional de Ciencias Naturales, c/o Jose Gutierrez Abascal No. 2, 28006 Madrid, Spain

Introduction

The anticipation of changes in environmental conditions may be considered as one of the principal mechanisms which explain adaptations of reproductive strategies (Baharav, 1983). The relation between environmental factors and the regulatory mechanisms of the reproductive cycle have been studied by different authors, both for animals in general (Lack, 1954; Gadil & Bossert, 1970; Schaffer, 1974) and ungulates in particular (Jarman, 1974; Leuthold & Leuthold, 1975; Nichols, 1978; Anderson, 1979; Baharav, 1983).

Little is known about reproduction in the Spanish ibex (Capra pyrenaica) (Couturier, 1962; Fandos, 1986). Asdell (1964) described some patterns of the genus Capra, but not of pyrenaica. The long gestation period of the Spanish ibex, the unpredictable variations of climate and availability of food, which are a consequence of the mediterranean climate, involve a high risk of mortality as much for the female goat as for the kid, especially if the latter is born when there is little food available or the weather conditions are adverse (Geist, 1971).

In the present work, the seasonality of reproduction, age of sexual maturity, length of gestation and litter size of the Spanish ibex are examined in different environmental and biological conditions

Methodology

This study compares the reproduction of two populations of the Spanish ibex in the south-east of the Iberian Peninsula. The main population lives in natural conditions occupying an area of 43,000 ha with a density of 6.2 ibex km⁻². The second population occupies an area of 64 ha in conditions of semi-captivity and is comprised of 92 ibex receiving a daily food supplement.

Direct observations for several days on female ibex were made monthly from October 1983 to September 1984, except during the times of birth and rut when sampling was carried out daily. Since individuals were not marked, results were based on the frequency of births (Caughley, 1977). The August sub-sample was chosen as the more representative, because by that month females have already given birth and mortality was still minimal. In the semi-captive population, the majority of the ibex gathered daily to take food so that mothers with offspring could be identified.

The age of the animals was determined by the number of segments and shape of horn, as described by Fandos (1986, In press) and Fandos et al., (In press).

TABLE I

Mean reproductive patterns of females of the two populations of Spanish ibex

Item Conception date	Sector of population of C. pyre- naica				
	Free-roaming	Semi-captivity			
	December	December			
Gestation period	155 days	155 days			
Parturition date	May	May			
Age of 1st conception	30 months	18 months			
Max. fawns at birth	1	2			
Natality rate	0.48	0.66			

Natality rate was defined as the product of the number of offspring and the number of two-yearold females in both free-roaming and semi-captive populations. The number of kids per female ibex was determined by which kids a female suckled.

The times of the year for copulation and parturition were determined in both populations from the first and last observations of their occurrence.

A known age sample of 15 foetuses was used for comparison of the time of parturition (Fandos, 1987).

Results

The reproductive patterns of the female Spanish ibex are given in Table I. The timing of parturition in both populations is fairly restricted, towards the end of spring to early summer. In semi-captivity this has been determined from the numbers of births taking place daily. In the case of free-roaming ibex, a 15-foetus sample has been used (Fig. 1). Rut occurs at the beginning of winter.

Sexual maturation of the females

The age of sexual maturity of the female ibex was different in the two populations. In semicaptivity there is a tendency to mature sexually at 18 months. Two of the nine females that reached two years of age in the 1984 sample each gave birth to a kid. In the free-roaming population, no two-year-old females were observed with offspring. This would suggest that, under normal conditions, the females tend to delay the age of maturity to at least 30 months.

Litter size

The number of offspring per pregnancy is dependent on genetic and environmental factors, the latter generally being the limiting one. In semi-captivity, 18 births were observed, of which four




FIG. 1. Frequency of parturition in 10-day intervals in free-roaming, ■; and semi-captive, □ ibex. The median parturition date is arrowed.

Table II
Numbers of females (fx) and young (Bx) in known age-classes
of the two populations of Spanish ibex

Age-class	Free-roaming			Semi-captivity		
	fx	Bx	Bx/fx	fx	Вх	Bx/fx
2 years old	6	0	0	9	2	0.22
3 years old	5	2	0.4	5	4	0.80
4-6 years old	13	10	0.76	5	5	1
7-11 years old	13	7	0.52	6	.6	1
12-14 years old	2	1	0.5	6	4	0.66
14 years and older	2	0	0	2	1	0.50

were twins (Table II). In this population, there was no scarcity of food since daily nourishment was provided. However, in the free-roaming population, 236 females were counted with one offspring and none with two. Similarly, in the survey of females shot during gestation period, none was observed with more than one foetus.

Discussion

The two study populations of ibex were separated more than 16 years ago. Both can be said to follow the k-strategy (low rate of population increase, advanced age of first reproduction, small litter size, repetition of the reproductive cycle, large body weight) (MacArthur & Wilson, 1967; Pianka, 1970).

Although the semi-captive population has an unlimited food supply throughout the year, the seasonality of reproduction is unaffected. The severity of winter, with the risk of births taking place in temperatures colder than -10 °C, could be the factor determining seasonality instead of the quality and quantity of food. No significant difference was observed between the two samples in respect of the combined median dates of birth (May 13) ($\chi^2 = 2 \cdot 1$; F.L. = 1; $P > 0 \cdot 1$) (Fig. 1).

The greater availability of food is reflected in the increased birth rate in the semi-captive population (larger litter size and earlier age of sexual maturity). The same effect has been observed in the population of duiker (Silvicapra) in Zambia when density and the intraspecific competition are diminished (Wilson & Roth in Delany, 1982).

To approach parturition with a shortage of food and without sufficient fat reserves in habitats subject to great variation involves a high risk to females owing to the energy demand of the young. Sexual maturity generally occurs in Spanish ibex after 30 months of age when their weight exceeds 24 kg. (Fandos, 1986). This relationship between the mother's weight and the onset of reproduction has been studied by several authors (Mitchell & Brown, 1974; Ellenberg, 1978; Saether & Haagenrud, 1983) who considered nutrition as one of the factors determining body size in ungulates (Verme, 1969; Kein, 1970). In the semi-captive population, food cannot be considered as a limiting factor and the critical weight of 24 kg will be reached before about 18 months, explaining births by two-year-old mothers. Woodgerd (1964) and Berger (1982) have also found a relationship between the mother's body weight and the age of first reproduction.

Litter size depends on genetic and environmental factors (Land, 1978); so the principal adaptation of Spanish ibex relates to the capacity of certain females, who are capable genetically of producing twins, to do so only when they have, or foresee the availability of, food in quantity and

quality. Thus the observed differences in reproductive performance can be explained either by the availability of food or the severity of winter.

REFERENCES

Asdell, S. A. (1964). Patterns of mammalian reproduction. New York: Cornell University Press.

Anderson, J. L. (1979). Reproductive seasonality of the nyala Tragelaphus angasi; the interaction of light, vegetation phenology, feeding style and reproductive physiology. Mamm. Rev. 9: 33-46.

Baharay, D. (1983). Reproductive strategies in female Mountain and Dorcas gazelles (Gazella gazella gazella and Gazella dorcas), J. Zool., Lond, 200: 445-453.

Berger, J. (1982). Female breeding age and lamb survival in desert bighorn sheep (Ovis canadensis). Mammalia 46: 183-190. Caughley, G. (1977). Analysis of vertebrate populations. London: John Wiley.

Couturier, M. (Ed.) (1962). Le bouquetin des Alpes (Capra aegagrus ibex ibex L.). Grenoble.

Delany, M. J. (1982). Mammal ecology. East Kilbride: Blackie & Son.

Ellenberg, H. (1978), Zur Populationsökologie des Rehes (Capreolus Capreolus L., Cervidae) in Mitteleuropa. Spixiana 1978 (Suppl. 2): 5-211.

Fandos, P. (1986). Aspectos ecológicos de la población de cabra montés en las Sierras de Cazorla y Segura, Jaen. PhD thesis, Universidad Complutense, Madrid.

Fandos, (1987). Croissance et developpement foetale du bouquetin de Cazorla. Mammalia 51: 579-585.

Fandos, P. (In press). Age determination in the Spanish mountain goat. Proc. XVIIIth Congr. Int. Un. Game Biologist 1987. Krakow.

Fandos, P., Vigal, C. R., Machordom, A., Fernandez, J. M., Martinez, T. & Palacios, F. (In press). Horn segments and dental cementun lines as age criteria for the Spanish Wild Goat, Capra pyrenaica Schinz, 1838. Säugetierk. Mitt.

Gadil, M. & Bossert, W. H. (1970). Life historical consequences of natural selection Am. Nat. 104: 1-24.

Geist, V. (1971). Mountain sheep: A study in behavior and evolution. Chicago: University of Chicago Press.

Jarman, R. J. (1974). The social organization of antelope in relation to their ecology. Behaviour 48: 215-267.

Kein, D. R. (1970). Food selection by North American deer and their response to over-utilization of preferred plant species. In Animal populations in relation to their food resources. Watson, A. (Ed.). Oxford: Blackwell.

Lack, D. (1954). The natural regulation of animal numbers. London: Oxford University Press.

Land, R. B. (1978). Reproduction in young sheep: some genetic and environmental sources of variation. J. Reprod. Fert. 52: 427-436.

Leuthold, W. & Leuthold, B. U. (1975). Temporal patterns of reproduction in ungulates in Tsavo East. E. Afr. Wildl. J. 13: 159-169.

MacArthur, R. H. & Wilson, E. O. (1967). The theory of island biogeography. Princeton, N J: Princeton University Press. Mitchell, B. & Brown, D. (1974). The effects of age and body size on fertility in female Red deer (Cervus elaphus L.). Proc. XIth Int. Congr. Game Biologist S.N.B., Stockholm.

Nichols, L. (1978). Dall sheep reproduction. J. Wildl. Mgmt 42: 570-580.

Pianka, E. R. (1970). On r and K selection. Am. Nat. 104: 592-597.

Saether, B.-E. & Haagenrud, H. (1983). Life history of the moose (Alces alces): Fecundity rates in relation to age and carcass weight. J. Mammal. 64: 226-232.

Schaffer, W. M. (1974). Optimal reproductive effort in fluctuating environments. Am. Nat. 108: 783-790.

Verme, L. J. (1969). Reproductive patterns of white-tailed deer related to nutritional plane. J. Wildl. Mgmt 33: 881-887.

Woodgerd, W. (1964). Population dynamics of bighorn sheep on Wildhorse Island. J. Wildl. Mgmt 28: 381-391.